

## Ankle Sprain



- Stay off your feet for 24 hours. Then walk slowly on the injured ankle.
- Use crutches or a cane until you can stand on your ankle without having pain.

Call your doctor **right away** if your:

- Bruising, swelling or pain gets worse
- Toes below the injury feel cold when you touch them, are numb or blue

**Talk to your doctor or nurse if you have any questions or concerns.**

The ankle is a joint where the bones of the legs and the feet meet. These bones are kept in place by ligaments. A sprain occurs when the ligaments are suddenly stretched or torn.

### Signs of an ankle sprain:

- Pain or tenderness
- Swelling
- Bruising
- Not able to move the ankle well

### Care of an ankle sprain:

Your doctor may want you to:

- Have an x-ray.
- Wear a splint or an ace wrap to support the injured ankle.
- Take over-the-counter pain medicine.
- Keep your foot raised above the level of your heart for the first days. This will reduce swelling and pain.
- Put ice on the injured ankle for 15-20 minutes each hour for 1-2 days. Put the ice in a plastic bag and place a towel between the bag of ice and your skin.

Text adapted from materials produced in 2006 by Health Information Translations ([www.healthinfotranslations.com](http://www.healthinfotranslations.com)), a partnership of The Ohio State University Medical Center, Mount Carmel Health and OhioHealth, Columbus, Ohio. This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivs License <http://creativecommons.org/licenses/by-nc-nd/3.0/us>

**A Healthy Roads Media project**  
[www.healthyroadsmedia.org](http://www.healthyroadsmedia.org)